

Warm Gas and Ionizing Photons in the Local Group

Philip R. Maloney

Center for Astrophysics and Space Astronomy, University of Colorado

J. Bland-Hawthorn

Anglo-Australian Observatory

Abstract. We investigate how much warm ($T \sim 10^6$ K) gas may exist in a Local Group corona, and, in particular, whether such a corona could be detected indirectly through its ionizing radiation. We calculate the non-LTE cooling of warm, low-density gas and the resulting ionizing photon emissivities for a range of metallicities. In principle a corona can produce a photon flux which significantly exceeds the cosmic ionizing background. However, the additional constraints which can be imposed on intragroup gas rule out models which produce detectable ionizing photon fluxes at the Milky Way's distance from the Local Group barycenter.

1. Introduction

How much warm ($T \sim 10^6$ K) gas could there be in the Local Group? This question dates back 40 years, to the classic Kahn & Woltjer (1959) “timing mass” paper; Kahn & Woltjer suggested that the bulk of the timing mass (which exceeded the visible stellar mass in the Local Group by a large factor) was in the form of a warm ($T \simeq 10^6$ K), low-density ($n \sim 10^{-4}$ cm $^{-3}$) plasma. This identification was driven largely by the difficulty of observing such gas directly. Although the mass of the Local Group is now believed to be dominated by (possibly non-baryonic) dark matter, the question of whether there is a substantial amount of warm gas present in the Local Group has recurred in several different contexts. In the following section we review the most recent suggestions for the presence of such a Local Group “corona”; in §3 we investigate whether such a corona could be detected indirectly through its ionizing radiation. A fuller discussion of this material is presented in Maloney & Bland-Hawthorn (1998a).

2. A Local Group Corona?

Suto *et al.* (1996) suggested that the measurement of the *COBE* cosmic microwave background (CMB) quadrupole moment could be significantly affected by the presence of a warm intragroup medium centered on the barycenter of the Local Group: for an electron density distribution of the form

$$n_e(r) = n_o \frac{r_o^2}{r^2 + r_o^2}, \quad (1)$$

where n_o is the central density and r_o is the core radius, centered a distance R from the Galaxy, the expected monopole and quadrupole anisotropies are

$$T_{0,sz} = \pi \theta_o \sigma_T \frac{kT}{mc^2} \frac{n_o r_o^2}{R}, \quad (2)$$

$$T_{2,sz} = \frac{\sqrt{5}\pi}{4} \left[\theta_o - 3 \frac{r_o}{R} + 3 \left(\frac{r_o}{R} \right)^2 \theta_o \right] \sigma_T \frac{kT}{mc^2} \frac{n_o r_o^2}{R}, \quad (3)$$

where $\theta_o \equiv \tan^{-1}(R/r_o)$. These equations can be inverted to get

$$n_o r_o T_{\text{keV}} < 7.4 \times 10^{21} \theta_o^{-1} \frac{R}{r_o} \left(\frac{y}{1.5 \times 10^{-5}} \right) \text{ cm}^{-2}. \quad (4)$$

$$n_o r_o T_{\text{keV}} < 1.6 \times 10^{20} Q_{\mu K} \frac{R}{r_o} \left[\theta_o - 3 \left(\frac{r_o}{R} \right) + 3 \theta_o \left(\frac{r_o}{R} \right)^2 \right]^{-1} \text{ cm}^{-2} \quad (5)$$

where the upper limit to the Compton y -parameter is $|y| = T_{0,sz}/2 < 1.5 \times 10^{-5}$ (95% confidence level: Fixsen *et al.* 1996), while the rms quadrupole amplitude $Q_{\text{RMS}} = 10^{-6} Q_{\mu K}$ K; the observed value $Q_{\mu K} \approx 6$ (*e.g.*, Bennett *et al.* 1996). Suto *et al.* argued that a LG corona which satisfied the inequality (4) could still significantly impact the CMB quadrupole measurement.

This conclusion was immediately contradicted by Banday & Gorski (1996), who demonstrated that the *COBE* data show no evidence for such a distortion, and by Pildis & McGaugh (1996), who pointed out that X-ray observations of poor groups, when fitted with density profiles of the form (1), typically obey $n_o r_o T_{\text{keV}} \lesssim 3 \times 10^{20} \text{ cm}^{-2}$, with $T_{\text{keV}} \sim 1$. Furthermore, most *detected* groups are dominated by ellipticals; the upper limits to $n_o r_o T_{\text{keV}}$ for spiral-rich groups are usually an order of magnitude smaller. Thus, although the *COBE* constraints on a LG corona are quite weak, analogy with similar (indeed, usually richer) poor groups suggests that the LG is unlikely to possess a significant gaseous halo.

However, Mulchaey *et al.* (1996), in an X-ray study of poor groups, suggested that spiral-rich groups might be undetected not because of an absence of gas, but because the gas is too cold to detect: spiral-dominated groups tend to have lower velocity dispersions, implying virial temperatures of only $T_{\text{vir}} \sim 0.2$ keV. The soft X-ray emission from gas at such temperatures is extremely difficult to detect. If the product $n_o r_o T_{\text{keV}}$ is similar to that in detected groups, the gas mass could be substantial: the mass within radius r is

$$M(r) \sim 7 \times 10^{11} \left(\frac{r_o}{100 \text{ kpc}} \right)^2 \left(\frac{r}{r_o} \right) \left(\frac{n_o r_o T_{\text{keV}}}{10^{20} \text{ cm}^{-2}} \right) \left(\frac{0.2}{T_{\text{keV}}} \right) M_{\odot} \quad (6)$$

for $r/r_o \gtrsim$ a few. (Note, however, that X-ray-detected poor groups usually have $M_{\text{gas}}/M_{\text{grav}} \sim 0.1$, which limits the total gas mass.) In fact, Wang & McCray (1993) found evidence from *ROSAT* observations for a diffuse thermal component, with $T_{\text{keV}} \sim 0.2$ and $n_e \sim 10^{-2} x^{-1/2} \text{ cm}^{-3}$, where x is the line-of-sight extent of the emitting region in kpc; the inferred electron density is smaller by a factor ~ 3 for solar metallicity rather than primordial gas. Since only the emission measure E_m is obtainable from the data, the spatial extent is unknown.

Blitz *et al.* (1998) have suggested that most HVCs are remnants of the formation of the Local Group, associated with continuing infall onto the LG. (See also Blitz, this volume.) If true, some fraction of these clouds will collide in the vicinity of the LG barycenter, and shock up to the virial temperature of the LG, $T \sim 2 \times 10^6$ K, leading to the formation of a warm intragroup medium.

In the next section we consider whether the ionizing photon flux from such a corona could be detectable, and the additional constraints which can at present be imposed on a LG halo.

3. Ionizing Photons from a Local Group Corona

We assume a density distribution of the form (1), with an outer boundary r_b and a temperature $T_{\text{kev}} \simeq 0.2$, the virial temperature of the LG. To calculate the emission and the ionizing photon flux from the gas, we have used the photoionization/shock code MAPPINGS, kindly provided by Ralph Sutherland. We consider metallicities $Z = 0.01, 0.1$, and $0.3 Z_\odot$. As long as $r_b \gtrsim$ a few r_o , the ionizing photon flux normally incident on a plane-parallel gas layer is

$$\phi_i(r) \approx 10^4 n_{-3}^2 r_{100} \left(\frac{\xi_i}{10^{-14}} \right) \frac{[0.8 + 1.3(r/r_o)^{1.35}]}{(1 + r^2/r_o^2)^{1.5}} \text{ cm}^{-2} \text{ s}^{-1} \quad (7)$$

where the central density $n_o = 10^{-3} n_{-3} \text{ cm}^{-3}$ and the core radius $r_o = 100 r_{100}$ kpc. For the temperatures and metallicities of interest, the integrated photon emissivity ξ_i generally lies between $3 \times 10^{-15} - 3 \times 10^{-14} \text{ cm}^{-3} \text{ s}^{-1} \text{ sr}^{-1}$. Poor groups show a very broad range of core radii, from tens to hundreds of kpc (Mulchaey *et al.* 1996), and typical central densities $n_o \sim$ a few $\times 10^{-3} \text{ cm}^{-3}$ (Pildis & McGaugh 1996). For these parameters, the photon fluxes predicted by equation (7) are quite substantial. (For reference, the cosmic ionizing background [the integrated contribution from AGN and galaxies] is probably $\phi_{i,\text{cos}} \sim 10^4 \text{ phot cm}^{-2} \text{ s}^{-1}$: Maloney & Bland-Hawthorn 1998b.) In Figure 1, we show (shaded in gray) the range of corona parameters (r_o, n_o) for which the resulting ionizing photon flux is between $\phi_i = 10^4$ and $\phi_i = 10^5 \text{ phot cm}^{-2} \text{ s}^{-1}$, for radial offsets $r = 0$ (upper region) and $r = 350 \text{ kpc}$ (lower region). The latter is the Galaxy's assumed radial distance r_{MW} from the center of the LG. However, there are additional constraints which we can place on a LG corona, which rule out a significant (*i.e.*, $\phi_i > \phi_{i,\text{cos}}$) contribution to the ionizing photon flux at r_{MW} :

- The assumption that any LG intragroup medium is typical of poor groups constrains the product $n_o r_o \lesssim 1.5 \times 10^{21} \text{ cm}^{-2}$, for an assumed temperature of $T_{\text{kev}} \approx 0.2$. This is plotted as the short-dashed line in Figure 1. Any corona which is not unusually rich must lie to the left of this line. This requirement alone rules out any significant contribution to ϕ_i at r_{MW} .
- The timing mass – obviously, the mass of the corona cannot exceed the mass of the LG. There are still come uncertainties in the determination of the timing mass (see Zaritsky 1994, and this volume, for a summary). We have adopted a value for the timing mass of $M_T = 5 \times 10^{12} M_\odot$ within $r = 1 \text{ Mpc}$ of the LG center. The mass in the corona is given by

$$M_c(r) = 4.3 \times 10^{11} n_{-3}^3 r_{100}^3 \left[1 - \frac{1}{x} \tan^{-1}(x) \right] M_\odot \quad (8)$$

where $x = r/r_o$, and for Figure 1 we have calculated the corona mass out to $r = 1$ Mpc. The timing mass constraint is shown as the solid line in Figure 1. As plotted, this is barely more restrictive than the *COBE* quadrupole limit (shown as the long-dashed line) and is only more restrictive than the assumption of a typical intragroup medium for large core radius. However, the true timing mass limit is considerably more stringent than this, as we must subtract off the masses of M31 and the Milky Way.

- Limits on the actual electron density at $r \sim r_{\text{MW}}$. These constraints come from two sources. First, observations of dispersion measures toward pulsars in the LMC and a distant globular cluster, NGC 5024 (Taylor, Manchester & Lyne 1993) require a mean $n_{-3} \sim 1$. However, most of this column must be contributed by the Reynolds layer, so only a small fraction (probably $\lesssim 10\%$) can be due to a LG corona. Secondly, an average density of no more than $n_{-3} \sim 0.1$ is allowed by models of the Magellanic Stream; otherwise, the Stream clouds would be plunging nearly radially into the Galaxy (Moore & Davis 1994). The hatched region in Figure 1 indicates the portion of (r_o, n_o) space in which $n_e(r_{\text{MW}}) \leq 10^{-4} \text{ cm}^{-3}$.




Figure 1. Constraints on a Local Group corona in the (r_o, n_o) plane. Coronae within the gray-shaded regions produce ionizing photon fluxes between $\phi_i = 10^5$ and $10^4 \text{ phot cm}^{-2} \text{ s}^{-1}$ (upper and lower edges) at radii $r = 0$ (lower region) and $r = 350 \text{ kpc}$ (upper region) with respect to the LG center. The long-dashed line is the *COBE* quadrupole constraint, the short-dashed line assumes the LG medium is “typical”, the solid line is the timing mass constraint, and the hatched region satisfies $n_e \leq 10^{-4} \text{ cm}^{-3}$ at $r = 350 \text{ kpc}$. See text for discussion.

The most stringent constraints come from the limits on the local value of n_e and from the timing mass (when realistic contributions from the Galaxy and M31 are included). Although it is possible to circumvent these restrictions to some extent by assuming the gas is highly clumpy, it is not possible to evade the soft X-ray determinations from Wang & McCray (1993) in this fashion, since this is a measurement of the emission measure. If the $T_{\text{kev}} \approx 0.2$ component inferred by Wang & McCray is extended on the scale of the local group, then the mean density is at most $n_{-3} \sim 0.3$, not far above the hatched region in Figure 1. These constraints rule out any significant ionizing photon flux at $r \sim r_{\text{MW}}$ from a Local Group corona: if the core density n_o is high, then r_o must be small, while if r_o is large, n_o must be low. LG coronae in the allowed region of (r_o, n_o) -space could produce ionizing fluxes much larger than the cosmic background, but only on scales of a few tens of kpc at best. An allowed LG medium could still contain a substantial amount of mass; the direct limits on such a medium have yet to improve on the values suggested by Kahn & Woltjer (1959).

Acknowledgments. PRM acknowledges support from the Astrophysical Theory Program through NASA grant NAG5-4061. We are grateful to Simon White for comments and platypus spotting.

References

- Bandy, A.J., & Górski, K.M. 1996, MNRAS, 283, L21
 Bennett C.L., *et al.* 1996, ApJ, 464, L1
 Blitz, L., Spergel, D.N., Teuben, P.J., Hartmann, D., & Burton, W.B. 1998, ApJ, in press.
 Fixsen, D.J., Cheng, E.S., Gales, J.M., Mather, J.C., Shafer, R.A., & Wright, E.L. 1996, ApJ, 473, 576
 Kahn, F.D., & Woltjer, L. 1959, ApJ, 130, 705
 Maloney, P.J., & Bland-Hawthorn, R.B. 1998a, ApJ, submitted.
 Maloney, P.J., & Bland-Hawthorn, R.B. 1998, in The Stromlo Workshop on High Velocity Clouds, ed. B.K. Gibson & M.E. Putnam (San Francisco: ASP), in press.
 Moore, B., & Davis, M. 1994, MNRAS, 270, 209
 Mulchaey, J.S., Davis, D.S., Mushotzky, R.F., & Burstein, D. 1996, ApJ, 456, 80
 Pildis, R.A., & McGaugh, S.S. 1996, ApJ, 470, L77
 Suto, Y., Makishima, K., Ishisaki, Y., & Ogasaka, Y. 1996, ApJ, 461, L33
 Taylor, J.H., Manchester, R.N., & Lyne, A.G. 1993, ApJS, 88, 529
 Wang, Q.D., & McCray, R.M. 1993, ApJ, 409, L37
 Zaritsky, D. 1994, in The Local Group: Comparative and Global Properties, 3rd CTIO/ESO Workshop, ed. A. Layden, R.C. Smith, & J. Storm (Garching: ESO), p. 187